

Carbohydrate & Insulin Adjustments

Common challenges of being active with diabetes

- Too little carbohydrate or too much insulin → low glucose (**hypoglycemia**)
- Too much carbohydrate or not enough insulin → high glucose (**hyperglycemia**)

Here are 4 strategies to manage glucose with physical activity

1. Modify carbohydrate intake before/during/after activity
2. Modify your meal-time bolus insulin
3. Set a temporary basal rate (if using an insulin pump)
4. Consider the type of activity

1. Modify Carbohydrate Intake Before/During/After Activity

- If you are adjusting carbohydrate intake only, aim to eat **1g of carb per kg per hour**. It is best to spread this over the course of the activity.
- Before or during activity, choose high glycemic index foods.
- If you have insulin on board or glucose is trending low, have a carbohydrate snack prior to activity without insulin.

2. Modify Your Meal-Time Bolus Insulin

- This is a good strategy to use if you plan to be active **within** 2 hours of a meal.
- Reducing meal-time insulin before activity can prevent low glucose. Try reducing by 25% for light activity, 50% for moderate activity, and 75% for high intensity activity.
- Reducing meal-time insulin following activity by 25-50% can prevent low glucose afterwards.
- Use caution when giving correction doses of insulin following activity. Consider reducing the correction dose by 50%.

3. Set a Temporary Basal Rate (If you use an Insulin Pump)

- This is a good strategy to use if you plan to be active **more than** 2 hours after a meal.
- For moderate to intense activity, start by reducing basal rate by 30-50% 1 hour before the start of activity until 2 hours after the activity.
- Note: If you take both mealtime and basal insulin and will be physically active for long durations, try reducing your daily basal insulin by 20% in addition to reducing your mealtime bolus before activity.

4. Consider the Type of Activity

- **Aerobic activity or 'cardio'** tends to lower glucose (ex: walking, jogging, bicycling). Risk of low glucose is higher for 24 hours following activity.
- **Anaerobic or 'resistance' activity** tends to raise glucose (ex: weight training, resistance bands, sprinting, high intensity interval training).
- **Competitive activities and team sports:** Practice/trainings days tend to lower glucose, while competition days tend to raise glucose. You might find that you need less of an insulin reduction on competition days.
- Tips to prevent low glucose
 - Performing resistance activity immediately before aerobic activity rather than performing aerobic activity alone or followed by resistance activity.
 - Perform brief (10 seconds), intermittent, maximal-intensity sprints either at the beginning, end or throughout a moderate-intensity activity session.

How Often Should I Check My Glucose When Active?

Activity can interfere with your ability to sense high or low glucose levels. It is recommended to check your glucose:

- 30 and 60 minutes before activity to determine direction of change
- Every 30 minutes during activity
- Within 30 minutes after the activity
- Before bed

Do Not Perform Physical Activity If

- You have had a severe low (<2.8mmol/L) within the last 24 hours
- Glucose is <4mmol/L; treat with quick sugar
- Glucose is >14.0mmol/L with ketones or >16.7mmol/L
 - Type 2 diabetes: Ensure proper hydration, monitor for signs and symptoms of dehydration.
 - Type 1 diabetes: Check for ketones. If ketones are 0.6mmol/L or greater, postpone activity and take a correction dose of insulin.

Putting It All Together

Many people find that a combination of these 4 strategies work best. The table below provides some ideas for combining methods.

	Anaerobic/ resistance activity	Aerobic/ cardio activity	Mixed activity
1-2 hrs before	<u>Correction bolus:</u> Consider if BG is trending high (ex: >9.0mmol/L) Temp basal: May not be needed	<u>Correction bolus:</u> Not needed unless glucose is high (>14.0mmol/L with ketones or >16.7mmol/L) Food bolus: ↓ by 25-75% Temp basal: ↓ rate by 25-50%	<u>Correction bolus:</u> Not needed unless glucose is high (>14.0mmol/L with ketones or >16.7mmol/L) Temp basal: ↓ rate by 25-50%
15-30 min before	Glucose <5.0mmol/L: Have 15g carb	Glucose <8.0mmol/L: Have 15g carb	
During	If activity is >30 minutes: Have 15-30g carbs per hour of activity	Have 0.5g carb/min of activity (ex: 15g carbs for 30 min of activity)	
After	Have a snack or meal that includes carb & protein <u>Food bolus:</u> usual dose <u>Correction dose:</u> ↓ by 25-50% Resume normal basal rates unless glucose <4mmol/L	Have a snack or meal that includes carb & protein <u>Correction and/or Food bolus:</u> ↓by 25%-50	

Adapted from: Riddell, Michael. Getting Pumped! An Insulin Pump Guide for Active Individuals with Type 1 Diabetes. Glue Inc. 2016

Other resources:

- www.excarbs.com
- www.diabetes.ca
- Walsh, John and Ruth Roberts. Pumping Insulin. Everything you Need to Succeed on an Insulin Pump.
- Scheiner, Gary. Think Like a Pancreas. A Practical Guide to Managing Diabetes with Insulin.

CARING FOR THE BODY, MIND & SPIRIT SINCE 1869

Renowned for compassionate care, St. Joseph's is one of the best academic health care organizations in Canada dedicated to helping people live to their fullest by minimizing the effects of injury, disease and disability through excellence in care, teaching and research.



sjhc.london.on.ca